arXiv:1509.06156v1 [math.AP] 21 Sep 2015

# DECOMPOSITIONS FOR HYPERGEOMETRIC FUNCTION $H_A(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z)$ , $H_B(\alpha,\beta_1,\beta_2;\gamma_1,\gamma,\gamma;x,y,z)$ , $H_C(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z)$

**Anvar H. Hasanov, Rakhila B. Seilkhanova, Roza D. Seilova**

S. Baishev University Aktobe, Department for Information Systems and Applied Mathematics,
030000 Aktobe, Zhubanov str. 302, Kazakhstan
E-Mails: anvarhasanov@yahoo.com, srahila@inbox.ru, roza_seilova@mail.ru

**Abstract.** With the help of some techniques based on certain inverse pairs of symbolic operators, the authors investigated several decomposition formulas associated with Srivastava's Hypergeometric functions of three variables. Some operator identities have been constructed in this matter. With the help these operator forms 15 decompositions are found which are expressed through product of Hypergeometric Gauss and Appell's functions.

**2000 Mathematics Subject Classification.** Primary 33C20, 33C65; Secondary 44A45.

**Key Words and Phrases.** Srivastava's hypergeometric functions, multiple hypergeometric functions, inverse pairs of symbolic operators, generalized hypergeometric function, integral representations.

## 1. Introduction.

A great interest to the theory of Hypergeometric functions of several variables was due to Solutions of many applied problems of differential equations of partial derivatives which are used with the help of such Hypergeometric functions. For instance, the energy absorbed with nonferromagnet conductor sphere included in internal magnet field, are calculated with the help of such functions [1]. The Hypergeometric functions of several variables are used in quantized chemistry as well [2]. Especially, in gas dynamics many problems lead to solution of degenerating second order partial differential equations. For instance, the problem of adiabatic flat-parallel without whirlwind gas flow and the flow problem of supersonic current from vessel with flat walls, also a lot of problems connecting with gas flow [3].

We note that the Riemann's functions and fundamental solutions of the degenerating second order partial differential equations are being defined by Hypergeometric functions of several variables [4]. In investigation the boundary value problems for these equations we need decompositions for Hypergeometric functions of several variables which are expressed through simple Hypergeometric Gauss and Appell's functions.

The operator method has been used in papers [5-7] and found decompositions for Hypergeometric functions of two variables which are expressed through Gauss Hypergeometric functions of one variables.

Over four decades ago by H.M.Srivastava was introduced Hypergeometric functions of three variables [8]

$$H_A\left(\alpha, \beta_1, \beta_2; \gamma_1, \gamma_2; x, y, z\right) = \sum_{m,n,p=0}^{\infty} \frac{(\alpha)_{m+p} (\beta_1)_{m+n} (\beta_2)_{n+p}}{(\gamma_1)_m (\gamma_2)_{n+p}\, m!n!p!} x^m y^n z^p, \tag{1.1}$$

$$H_B\left(\alpha, \beta_1, \beta_2; \gamma_1, \gamma_2, \gamma_3; x, y, z\right) = \sum_{m,n,p=0}^{\infty} \frac{(\alpha)_{m+p} (\beta_1)_{m+n} (\beta_2)_{n+p}}{(\gamma_1)_m (\gamma_2)_n (\gamma_3)_p\, m!n!p!} x^m y^n z^p, \tag{1.2}$$

$$H_C\left(\alpha, \beta_1, \beta_2; \gamma\, ; x, y, z\right) = \sum_{m,n,p=0}^{\infty} \frac{(\alpha)_{m+p} (\beta_1)_{m+n} (\beta_2)_{n+p}}{(\gamma)_{m+n+p}\, m!n!p!} x^m y^n z^p. \tag{1.3}$$

In this work we construct decompositions for Hypergeometric functions (1.1)-(1.2) with the help of Burchnall and Chaundy [5-7] method. These decompositions consist of simple Appell's functions and Gauss functions. Further, by means of obtained decompositions we determine some definite integrals connecting with functions (1.1)-(1.2).

## 2. Main operators.

Over six decades ago, Burchnall and Chaundy [5, 6] and Chaundy [7] systematically presented a number of expansion and decomposition formulas for double hypergeometric functions in series of simplest Hypergeometric functions. Their method is based on following inverse pairs of symbolic operators:

$$\nabla_{xy}(h) := \frac{\Gamma(h)\Gamma(\delta_1 + \delta_2 + h)}{\Gamma(\delta_1 + h)\Gamma(\delta_2 + h)} = \sum_{k=0}^{\infty} \frac{(-\delta_1)_k (-\delta_2)_k}{(h)_k\, k!}, \tag{2.1}$$

$$\Delta_{xy}(h) := \frac{\Gamma(\delta_1 + h)\Gamma(\delta_2 + h)}{\Gamma(h)\Gamma(\delta_1 + \delta_2 + h)} = \sum_{k=0}^{\infty} \frac{(-\delta_1)_k (-\delta_2)_k}{(1 - h - \delta_1 - \delta_2)_k\, k!}$$

$$= \sum_{k=0}^{\infty} \frac{(-1)^k (h)_{2k} (-\delta_1)_k (-\delta_2)_k}{(h+k-1)_k (\delta_1 + h)_k (\delta_2 + h)_k \, k!}, \tag{2.2}$$

$$\begin{aligned}
\nabla_{xy}(h)\,\Delta_{xy}(l) &:= \frac{\Gamma(h)\,\Gamma(\delta_1+\delta_2+h)}{\Gamma(\delta_1+h)\,\Gamma(\delta_2+h)} \frac{\Gamma(\delta_1+l)\,\Gamma(\delta_2+l)}{\Gamma(l)\,\Gamma(\delta_1+\delta_2+l)} \\
&= \sum_{k=0}^{\infty} \frac{(l-h)_k (l)_{2l} (-\delta_1)_k (-\delta_2)_k}{(l+k-1)_k (l+\delta_1)_k (l+\delta_2)_k \, k!} \\
&= \sum_{k=0}^{\infty} \frac{(h-l)_k (l)_{2l} (-\delta_1)_k (-\delta_2)_k}{(h)_k (1-l-\delta_1-\delta_2)_k \, k!}, \quad (\delta_1 := x\frac{\partial}{\partial x};\ \delta_2 := y\frac{\partial}{\partial y}).
\end{aligned} \tag{2.3}$$

We now introduce here the following multivariable analogues of the Burchnall- Chaundy symbolic operators $\nabla_{xy}(h)$ and $\Delta_{xy}(h)$ defined by (2.1) - (2.3), respectively (cf. [9, p. 240]; see [10, p. 113] for the case when $r = 3$ ):

$$\begin{aligned}
\tilde{\nabla}_{x_1;x_2\cdots x_r}(h) &:= \frac{\Gamma(h)\,\Gamma(\delta_1+\cdots+\delta_r+h)}{\Gamma(\delta_1+h)\,\Gamma(\delta_2+\cdots+\delta_r+h)} = \\
&= \sum_{k_2,\ldots,k_r=0}^{\infty} \frac{(-\delta_1)_{k_2+\ldots+k_r} (-\delta_2)_{k_2} \cdots (-\delta_r)_{k_r}}{(h)_{k_2+\ldots+k_r} \, k_2! \cdots k_r!},
\end{aligned} \tag{2.4}$$

and

$$\begin{aligned}
\tilde{\Delta}_{x_1;x_2\cdots x_r}(h) &:= \frac{\Gamma(\delta_1+h)\,\Gamma(\delta_2+\delta_3+\cdots+\delta_r+h)}{\Gamma(h)\,\Gamma(\delta_1+\delta_2+\cdots+\delta_r+h)} \\
&= \sum_{k_2,\ldots,k_r=0}^{\infty} \frac{(-\delta_1)_{k_2+\ldots+k_r} (-\delta_2)_{k_2} \cdots (-\delta_r)_{k_n}}{(1-h-\delta_1-\delta_2-\cdots-\delta_r)_{k_2+\ldots+k_r} \, k_2! \cdots k_r!} \\
&= \sum_{k_2,\ldots,k_r=0}^{\infty} \frac{(-1)^{k_2+\ldots+k_r} (\gamma)_{2(k_2+\ldots+k_r)} (-\delta_1)_{k_2+\ldots+k_r} (-\delta_2)_{k_2} \cdots (-\delta_n)_{k_r}}{(\gamma+k_2+\ldots+k_r-1)_{k_2+\ldots+k_r} (\delta_1+\gamma)_{k_2+\ldots+k_r} (\delta_2+\delta_3+\cdots+\delta_r+\gamma)_{k_2+\ldots+k_r} \, k_2! \cdots k_r!},
\end{aligned} \tag{2.5}$$

$$(\delta_j := x_j \frac{\partial}{\partial x_j};\ j = 1, \ldots, r),$$

where we have applied such known multiple hypergeometric summation as (cf. [11]; see also [12, p. 117])

$$F_D^{(r)}(a; b_1, \cdots, b_r; c; 1, \cdots, 1) = \frac{\Gamma(c)\,\Gamma(c-a-b_1-\cdots-b_r)}{\Gamma(c-a)\,\Gamma(c-b_1-\cdots-b_r)}, \tag{2.6}$$

$$(\mathrm{Re}\,(c-a-b_1-\cdots-b_r) > 0;\ c \notin Z_0 := \{0, -1, -2, -3, \cdots\}),$$

for the Lauricella function $F_D^{(r)}$ in $r$ variables, defined by (cf. [11]; see also [13, p.33, Equation 1.4 (4)])

$$F_D^{(r)}\left(a;b_1,\cdot\cdot\cdot,b_r;c;x_1,x_2,\cdot\cdot\cdot,x_r\right):=$$

$$:=\sum_{m_1,m_2,\cdots,m_r=0}^{\infty}\frac{(a)_{m_1+m_2+\cdots+m_r}(b_1)_{m_1}\cdots(b_r)_{m_r}}{(c)_{m_1+m_2+\cdots+m_r}m_1!m_2!\cdots m_r!}x_1^{m_1}x_2^{m_2}\cdots x_r^{m_r}, \tag{2.7}$$

$$\left(\max\left\{|x_1|,...,\ |x_r|\right\}<1\right),$$

$(h)_k$ - Pochhammer's symbol.

## 3. Main operator identities.

According to operators (2.1)-(2.5) for Hypergeometric functions (1.1)-(1.3) we find the following operator identities

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F_1\left(\beta_2;\beta_1,\alpha;\gamma_2;y,z\right), \tag{3.1}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma,\gamma;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\gamma\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F_1\left(\beta_2;\beta_1,\alpha;\gamma;y,z\right), \tag{3.2}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\tilde{\Delta}_x\left(\gamma\right)F\left(\alpha,\beta_1;\gamma\,;x\right)F_1\left(\beta_2;\beta_1,\alpha;\gamma\,;y,z\right), \tag{3.3}$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\Delta_{yz}\left(\gamma_2\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F_2\left(\beta_2;\beta_1,\alpha;\gamma_2,\gamma_2;y,z\right), \tag{3.4}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F_2\left(\beta_2;\beta_1,\alpha;\gamma_2,\gamma_3;y,z\right), \tag{3.5}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\gamma\right)\tilde{\Delta}_x\left(\gamma\right)F\left(\alpha,\beta_1;\gamma\,;x\right)F_2\left(\beta_2;\beta_1,\alpha;\gamma,\gamma;y,z\right), \tag{3.6}$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\beta_2\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F_3\left(\beta_1,\beta_2,\beta_2,\alpha;\gamma_2;y,z\right), \tag{3.7}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma,\gamma;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\beta_2\right)\nabla_{yz}\left(\gamma\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F_3\left(\beta_2,\alpha,\beta_1,\beta_2;\gamma;y,z\right), \tag{3.8}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\beta_2\right)\tilde{\Delta}_x\left(\gamma\right)F\left(\alpha,\beta_1;\gamma;x\right)F_3\left(\beta_2,\beta_2,\beta_1,\alpha;\gamma;y,z\right), \tag{3.9}$$

$$H_A\left(\alpha,\alpha,\beta_2;\gamma_1,\gamma_2;x,y,z\right)$$

$$=\nabla_{xy}\left(\alpha\right)\nabla_{xz}\left(\alpha\right)\Delta_{yz}\left(\alpha\right)\Delta_{yz}\left(\gamma_2\right)F\left(\alpha,\alpha;\gamma_1;x\right)F_4\left(\alpha,\ \beta_2;\gamma_2,\gamma_2;y,z\right), \tag{3.10}$$

$$H_B\left(\alpha,\alpha,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\alpha\right)\Delta_{yz}\left(\alpha\right)F\left(\alpha,\alpha;\gamma_1;x\right)F_4\left(\alpha,\beta_2,\gamma_2,\gamma_3;y,z\right), \tag{3.11}$$

$$H_C\left(\alpha,\alpha,\beta_2;\gamma\,;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\alpha\right)\Delta_{yz}\left(\alpha\right)\Delta_{yz}\left(\gamma\right)\tilde{\Delta}_x\left(\gamma\right)F\left(\alpha,\alpha;\gamma;x\right)F_4\left(\alpha,\beta_2,\gamma,\gamma;y,z\right), \tag{3.12}$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\beta_2\right)\Delta_{yz}\left(\gamma_2\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F\left(\beta_1,\beta_2;\gamma_2;y\right)F\left(\alpha,\beta_2;\gamma_2;z\right), \tag{3.13}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\beta_2\right)F\left(\alpha,\beta_1;\gamma_1;x\right)F\left(\beta_1,\beta_2;\gamma_2;y\right)F\left(\alpha,\beta_2;\gamma_3;z\right), \tag{3.14}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)$$

$$=\nabla_{xz}\left(\alpha\right)\nabla_{xy}\left(\beta_1\right)\nabla_{yz}\left(\beta_2\right)\Delta_{yz}\left(\gamma\right)\tilde{\Delta}_x\left(\gamma\right)F\left(\alpha,\beta_1;\gamma\,;x\right)F\left(\beta_1,\beta_2;\gamma\,;y\right)F\left(\alpha,\beta_2;\gamma\,;z\right), \tag{3.15}$$

The operator identities for Hypergeometric functions $H_A\,, H_B, H_C$ can be proved with the help of Mellin's transformation [14]. For instance, we show that the operator identity (3.1) is true. According to known E.W.Barnes definition [12, p.40] we have

$$F\left(a;b;c;x\right)=\frac{1}{2\pi i}\frac{\Gamma\left(c\right)}{\Gamma\left(a\right)\Gamma\left(b\right)}\int\limits_{-i\infty}^{+i\infty}\Gamma\left[\begin{array}{c}a+s_1,b+s_1,-s_1\\c+s_1\end{array}\right]\left(-x\right)^{s_1}ds_1, \tag{3.16}$$

$c\neq 0,-1,-2,...$ and $\mathrm{Re}\,s_1<0$, Re $\left(a+s_1\right)>0$, Re $\left(b+s_1\right)>0$ ,

$$F_1\left(a;b_1,b_2;c;x,y\right)=-\tfrac{1}{4\pi^2}\tfrac{\Gamma(c)}{\Gamma(a)\Gamma(b_1)\Gamma(b_2)}$$

$$\cdot\int\limits_{-i\infty}^{+i\infty}\int\limits_{-i\infty}^{+i\infty}\Gamma\left[\begin{array}{c}a+s_1+s_2,b_1+s_1,b_2+s_2,-s_1,-s_2\\c+s_1+s_2\end{array}\right]\left(-x\right)^{s_1}\left(-y\right)^{s_2}ds_1ds_2, \tag{3.17}$$

$c\neq 0,-1,-2,...$ and

$\mathrm{Re}\,s_1<0$, $\mathrm{Re}\,s_2<0$, Re $\left(a+s_1+s_2\right)>0$, Re $\left(b_1+s_1\right)>0$, Re $\left(b_2+s_2\right)>0$,

By using equality

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=\sum_{m=0}^{\infty}\frac{\left(\alpha\right)_m\left(\beta_1\right)_m}{\left(\gamma_1\right)_m m!}x^m F_1\left(\beta_2;\beta_1+m,\alpha+m;\gamma_2;y,z\right),$$

we find

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=-\frac{1}{8\pi^2 i}\frac{\Gamma\left(\gamma_1\right)\Gamma\left(\gamma_2\right)}{\Gamma\left(\alpha\right)\Gamma\left(\beta_1\right)\Gamma\left(\beta_2\right)}$$

$$\cdot\int\limits_{-i\infty}^{+i\infty}\int\limits_{-i\infty}^{+i\infty}\int\limits_{-i\infty}^{+i\infty}\Gamma\left[\begin{array}{c}\alpha+s_1+s_3,\beta_1+s_1+s_2,\beta_2+s_2+s_3,-s_1,-s_2,-s_3\\\gamma_1+s_1,\gamma_2+s_2+s_3\end{array}\right]$$

$$\cdot\left(-x\right)^{s_1}\left(-y\right)^{s_2}\left(-z\right)^{s_3}ds_1ds_2ds_3, \tag{3.18}$$

$\gamma_1$ , $\gamma_2$ $\neq$ $0,-1,-2,...$ and $\mathrm{Re}\,s_1<0$, $\mathrm{Re}\,s_2<0$, $\mathrm{Re}\,s_2<0$, Re $\left(\alpha+s_1+s_3\right)>0$, Re $\left(\beta_1+s_1+s_2\right)>0$, Re $\left(\beta_2+s_2+s_3\right)>0$, where

$$\Gamma\left[\begin{array}{c}a_1,a_2,...,a_m\\b_1,b_2,...,b_n\end{array}\right]=\frac{\Gamma\left(a_1\right)\Gamma\left(a_2\right)\cdots\Gamma\left(a_m\right)}{\Gamma\left(b_1\right)\Gamma\left(b_2\right)\cdots\Gamma\left(b_n\right)}.$$

Therefore, from equalities (3.16)-(3.18) we can have the following

$$F\left(a;b;c;x\right)\leftrightarrow\frac{\Gamma\left(c\right)}{\Gamma\left(a\right)\Gamma\left(b\right)}\Gamma\left[\begin{array}{c}a+s_1,b+s_1,-s_1\\c+s_1\end{array}\right], \tag{3.19}$$

$c\neq 0,-1,-2,...$ and $\mathrm{Re}\,s_1<0$, Re $\left(a+s_1\right)>0$, Re $\left(b+s_1\right)>0$ ,

$$F_1(a;b_1,b_2;c;x,y) \leftrightarrow \frac{\Gamma(c)}{\Gamma(a)\Gamma(b_1)\Gamma(b_2)}\Gamma\left[\begin{array}{c} a+s_1+s_2, b_1+s_1, b_2+s_2, -s_1, -s_2 \\ c+s_1+s_2 \end{array}\right], \tag{3.20}$$

$c \neq 0,-1,-2,...$ and $\operatorname{Re} s_1 < 0$, $\operatorname{Re} s_2 < 0$, $\operatorname{Re}(a+s_1+s_2) > 0$, $\operatorname{Re}(b_1+s_1) > 0$, $\operatorname{Re}(b_2+s_2) > 0$ ,

$$\begin{aligned} &H_A(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z) \\ &\leftrightarrow \frac{\Gamma(\gamma_1)\Gamma(\gamma_2)}{\Gamma(\alpha)\Gamma(\beta_1)\Gamma(\beta_2)}\Gamma\left[\begin{array}{c} \alpha+s_1+s_3, \beta_1+s_1+s_2, \beta_2+s_2+s_3, -s_1, -s_2, -s_3 \\ \gamma_1+s_1, \gamma_2+s_2+s_3 \end{array}\right], \end{aligned} \tag{3.21}$$

$\gamma_1$ , $\gamma_2 \neq 0,-1,-2,...$ and $\operatorname{Re} s_1 < 0, \operatorname{Re} s_2 < 0$, $\operatorname{Re} s_2 < 0$, $\operatorname{Re}(\alpha+s_1+s_3) > 0$, $\operatorname{Re}(\beta_1+s_1+s_2) > 0$, $\operatorname{Re}(\beta_2+s_2+s_3) > 0$ .

At present we proof identity (3.1). Applying Mellin's transformation in both parts of (3.1) and taking into account equality

$$(-\delta_i)_k f(x) \leftrightarrow (-s_i)_k f^*(s_i), \tag{3.22}$$

$i = 1,2,3$ , moreover $\lim\limits_{x\to 0} x^{s-j-1} f^{(j)}(x) = 0$, $j = 0,1,...,k-1$, and from definition for (2.1) we have

$$\begin{aligned} &H_A(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z) \\ &\leftrightarrow \frac{\Gamma(\gamma_1)\Gamma(\gamma_2)}{\Gamma^2(\alpha)\Gamma^2(\beta_1)\Gamma(\beta_2)}\Gamma\left[\begin{array}{c} \alpha+s_1, \alpha+s_3, \beta_1+s_1, \beta_1+s_2, \beta_2+s_2+s_3, -s_1, -s_2, -s_3, \\ \gamma_1+s_1, \gamma_2+s_2+s_3 \end{array}\right] \\ &\cdot \sum_{i,j=0}^{\infty} \frac{(-s_1)_i(-s_1)_j(-s_2)_j(-s_3)_i}{(\alpha)_i(\beta_1)_j\, i!j!} \end{aligned} \tag{3.23}$$

By virtue of identity [15],

$$F(a,b;c;1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)},$$
$$(\operatorname{Re}(c-a-b > 0,\ \ c \notin Z_0 := \{0,-1,-2,-3,\cdots\}),$$

for expression (3.23) we have the following

$$\sum_{i,j=0}^{\infty} \frac{(-s_1)_i(-s_1)_j(-s_2)_j(-s_3)_i}{(\alpha)_i(\beta_1)_j\, i!j!} = \frac{\Gamma(\alpha)\Gamma(\alpha+s_1+s_3)}{\Gamma(\alpha+s_1)\Gamma(\alpha+s_3)}\frac{\Gamma(\beta_1)\Gamma(\beta_1+s_1+s_2)}{\Gamma(\beta_1+s_1)\Gamma(\beta_1+s_2)}. \tag{3.24}$$

Substituting equalities (3.24) into expression (3.23) and after simplest calculations we derive the operator identities (3.1). Analogically, with the help of Mellin's transformation can be proved the other operator identities.

## 4. The Decompositions for Hypergeometric functions $H_A$, $H_B$, $H_C$.

Applying operators and superposition of operators from identities (3.1)-(3.15) we have the following

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=\sum_{i,j=0}^{\infty}\frac{(\alpha)_{i+j}\left(\beta_1\right)_{i+j}\left(\beta_2\right)_{i+j}}{\left(\gamma_1\right)_{i+j}\left(\gamma_2\right)_{i+j}\,i!j!}x^{i+j}y^{j}z^{i}\cdot$$

$$\cdot F\left(\alpha+i+j,\beta_1+i+j;\gamma_1+i+j;x\right)F_1\left(\beta_2+i+j;\beta_1+i+j,\alpha+i;\gamma_2+i+j;y,z\right), \tag{4.1}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma,\gamma;x,y,z\right)=\sum_{i,j,k=0}^{\infty}\frac{(\alpha)_{i+j+k}\left(\beta_1\right)_{i+j+k}\left(\beta_2\right)_{2i+j+k}}{\left(\gamma_1\right)_{j+k}\left(\gamma\right)_{i}\left(\gamma\right)_{2i+j+k}\,i!j!k!}x^{j+k}y^{i+j}z^{i+k}$$

$$F\left(\alpha+i+j+k,\beta_1+i+j+k;\gamma_1+j+k;x\right)$$

$$F_1\left(\beta_2+2i+j+k;\beta_1+i+j,\alpha+i+j+k;\gamma+2i+j+k;y,z\right), \tag{4.2}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)=$$

$$=\sum_{i,j,k,l=0}^{\infty}\frac{(-1)^{i+j}(\alpha)_{i+2j+k+l}\left(\beta_1\right)_{2i+j+k+l}\left(\beta_2\right)_{i+j+k+l}\left(\gamma\right)_{2i+2j}}{\left(\gamma+i+j-1\right)_{i+j}\left(\gamma\right)_{2i+2j+k+l}\left(\gamma\right)_{2i+2j+k+l}\,i!j!k!l!}x^{i+j+k+l}y^{i+k}z^{j+l}\cdot$$

$$\cdot F\left(\alpha+i+2j+k+l,\beta_1+2i+j+k+l;\gamma+2i+2j+k+l\,;x\right)$$

$$F_1\left(\beta_2+i+j+k+l;\beta_1+2i+j+k,\alpha+i+2j+k+l;\gamma+2i+2j+k+l\,;y,z\right), \tag{4.3}$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=$$

$$=\sum_{i,j,k=0}^{\infty}(-1)^k\frac{\left(\gamma_2\right)_{2k}(\alpha)_{i+j+k}\left(\beta_1\right)_{i+j+k}\left(\beta_2\right)_{i+j+2k}}{\left(\gamma_2+k-1\right)_{k}\left(\gamma_1\right)_{i+j}\left(\gamma_2\right)_{i+2k}\left(\gamma_2\right)_{j+2k}\,i!j!k!}x^{i+j}y^{j+k}z^{i+k}$$

$$F\left(\alpha+i+j+k,\beta_1+i+j+k;\gamma_1+i+j;x\right)$$

$$F_2\left(\beta_2+i+j+2k;\beta_1+j+k,\alpha+i+j+k;\gamma_2+j+2k,\gamma_2+i+2k;y,z\right), \tag{4.4}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)=\sum_{i,j=0}^{\infty}\frac{(\alpha)_{i+j}\,(\beta_1)_{i+j}\,(\beta_2)_{i+j}}{(\gamma_1)_{i+j}\,(\gamma_2)_j\,(\gamma_3)_i\,i!j!}x^{i+j}y^jz^i\cdot$$

$$\cdot F\left(\alpha+i+j,\beta_1+i+j;\gamma_1+i+j;x\right)F_2\left(\beta_2+i+j;\beta_1+i+j,\alpha+i;\gamma_2+j,\gamma_3+i;y,z\right) \tag{4.5}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)=$$

$$=\sum_{i,j,k,l,r=0}^{\infty}\frac{(-1)^{i+j+r}\,(\alpha)_{i+2j+k+l+r}\,(\beta_1)_{2i+j+k+l}\,(\beta_1)_{2i+j+k+r}\,(\beta_2)_{i+j+k+l+2r}}{(\gamma+i+j-1)_{i+j}\,(\gamma+2i+2j+k+l+r-1)_r\,(\beta_1)_{2i+j+k}}$$

$$\frac{(\gamma)_{2i+2j}}{(\gamma)_{2i+2j+k+l}\,(\gamma)_{2i+2j+k+l+2r}\,i!j!k!l!r!}x^{i+j+k+l}y^{i+k+r}z^{j+l+r}\cdot$$

$$\cdot F\left(\alpha+i+2j+k+l,\beta_1+2i+j+k+l;\gamma+2i+2j+k+l\,;x\right)\cdot$$

$$F_2(\beta_2+i+j+k+l+2r;\beta_1+2i+j+k+r,\alpha+i+2j+k+l+r;$$

$$\gamma+2i+2j+k+l+2r,\gamma+2i+2j+k+l+2r;y,z), \tag{4.6}$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=\sum_{i,j,k=0}^{\infty}\frac{(\alpha)_{i+j+k}\,(\beta_1)_{i+j+k}\,(\beta_2)_{i+j}\,(\beta_2)_{i+k}}{(\beta_2)_i\,(\gamma_1)_{j+k}\,(\gamma_2)_{2i+j+k}\,i!j!k!}x^{j+k}y^{i+j}z^{i+k}$$

$$F\left(\alpha+i+j+k,\beta_1+i+j+k;\gamma_1+j+k;x\right)$$

$$F_3\left(\beta_1+i+j,\beta_2+i+k,\beta_2+i+j,\alpha+i+j+k;\gamma_2+2i+j+k;y,z\right), \tag{4.7}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma,\gamma;x,y,z\right)=$$

$$=\sum_{i,j,k,l=0}^{\infty}\frac{(\alpha)_{j+2k+l}\,(\beta_1)_{i+j+k+l}\,(\beta_2)_{2i+j+k}\,(\beta_2)_{2i+j+l}}{(\beta_2)_{2i+j}\,(\gamma)_i\,(\gamma)_{2i+2j+k+l}\,(\gamma_1)_{k+l}\,i!j!k!l!}x^{k+l}y^{i+j+k}z^{i+j+l}\cdot$$

$$\cdot F\left(\alpha+j+2k+l,\beta_1+i+j+k+l;\gamma_1+k+l;x\right)\cdot$$

$$\cdot F_3(\beta_2+2i+j+k,\alpha+i+j+k+l,\beta_1+i+j+k,\beta_2+2i+j+l;$$

$$\gamma+2i+2j+k+l;y,z), \tag{4.8}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)=$$

$$\sum_{i,j,k,l,r=0}^{\infty}\frac{(-1)^{i+j}\,(\alpha)_{i+2j+k+l}\,(\alpha)_{i+2j+k+r}\,(\beta_1)_{2i+j+k+l+r}\,(\beta_2)_{i+j+k+l+r}\,(\gamma)_{2i+2j}}{(\gamma+i+j-1)_{i+j}\,(\alpha)_{i+2j+k}\,(\gamma)_{2i+2j+k+l}\,(\gamma)_{2i+2j+k+l+2r}\,i!j!k!l!r!}\cdot$$

$$\cdot x^{i+j+k+l}y^{i+l+r}z^{j+k+r}\cdot F\left(\alpha+i+2j+k+l,\beta_1+2i+j+k+l;\gamma+2i+2j+k+l;x\right)$$

$$F_3(\beta_2+i+j+k+l+r,\beta_2+i+j+k+l+r,\beta_1+2i+j+k+l+r,\alpha+i+2j+k+r;$$

$$;\gamma+2i+2j+k+l+2r;y,z), \tag{4.9}$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=$$

$$=\sum_{i,j,k=0}^{\infty}\frac{(\alpha)_{i+j}\,(\alpha)_{i+k}\,(\beta_1)_{i+j+k}\,(\beta_2)_{i+j+k}\,(\gamma_2-\beta_2)_k}{(\gamma_2+i+j+k-1)_k\,(\alpha)_i\,(\gamma_1)_{i+j}\,(\gamma_2)_{i+j+2k}\,i!j!k!}x^{i+j}y^{j+k}z^{i+k}\cdot$$

$$\cdot F\left(\alpha+i+j,\beta_1+i+j;\gamma_1+i+j;x\right)\cdot$$

$$\cdot F\left(\beta_2+i+j+k,\beta_1+i+j+k;\gamma_2+i+j+2k;y\right)\cdot$$

$$\cdot F\left(\beta_2+i+j+k,\alpha+i+k;\gamma_2+i+j+2k;z\right), \tag{4.10}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)=\sum_{i,j,k=0}^{\infty}\frac{(\alpha)_{i+j+k}\,(\beta_1)_{i+j+k}\,(\beta_2)_{i+j}\,(\beta_2)_{j+k}}{(\beta_2)_j\,(\gamma_1)_{i+k}\,(\gamma_2)_{i+j}\,(\gamma_3)_{j+k}\,i!j!k!}x^{i+k}y^{i+j}z^{j+k}\cdot$$

$$F\left(\alpha+i+j+k,\beta_1+i+j+k;\gamma_1+i+k;x\right)F\left(\beta_1+i+j,\beta_2+i+j;\gamma_2+i+j;y\right)$$

$$F\left(\beta_2+j+k,\alpha+i+j+k;\gamma_3+j+k;z\right), \tag{4.11}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,z\right)=$$

$$=\sum_{i,j,k,l,r=0}^{\infty}\frac{(-1)^{i+j}\,(\alpha)_{i+2j+k+l+r}\,(\beta_1)_{2i+j+k+l}\,(\beta_1)_{2i+j+k+r}\,(\beta_2)_{i+j+k+l+r}}{(\gamma+i+j-1)_{i+j}\,(\gamma+2i+2j+k+l+r-1)_r\,(\beta_1)_{2i+j+k}}$$

$$\frac{(\gamma)_{2i+2j}\,(\gamma-\beta_2)_{i+j+r}}{(\gamma-\beta_2)_{i+j}\,(\gamma)_{2i+2j+k+l}\,(\gamma)_{2i+2j+k+l+2r}\,i!j!k!l!r!}x^{i+j+k+l}y^{i+k+r}z^{j+l+r}\cdot$$

$$\cdot F\left(\alpha+i+2j+k+l,\beta_1+2i+j+k+l;\gamma+2i+2j+k+l\,;x\right)\cdot$$

$$\cdot F\left(\beta_2+i+j+k+l+r,\beta_1+2i+j+k+r;\gamma+2i+2j+k+l+2r;y\right)\cdot$$

$$\cdot F\left(\beta_2+i+j+k+l+r,\alpha+i+2j+k+l+r;\gamma+2i+2j+k+l+2r;z\right), \tag{4.12}$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,y\right)=\sum_{i,j=0}^{\infty}\frac{(\alpha)_{i+j}\,(\beta_1)_{i+j}\,(\beta_2)_{i+j}}{(\gamma_1)_{i+j}\,(\gamma_2)_{i+j}\,i!j!}x^{i+j}y^{i+j}\cdot$$

$$\cdot F\left(\alpha+i+j,\beta_1+i+j;\gamma_1+i+j;x\right)F\left(\beta_2+i+j;\alpha+\beta_1+2i+j;\gamma_2+i+j;y\right) \tag{4.13}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma,\gamma;x,y,y\right)=\sum_{i,j,k=0}^{\infty}\frac{(\alpha)_{i+j+k}\,(\beta_1)_{i+j+k}\,(\beta_2)_{2i+j+k}}{(\gamma_1)_{j+k}\,(\gamma)_i\,(\gamma)_{2i+j+k}\,i!j!k!}x^{j+k}y^{2i+j+k}$$

$$F\left(\alpha+i+j+k,\beta_1+i+j+k;\gamma_1+j+k;x\right)$$

$$F\left(\beta_2+2i+j+k;\alpha+\beta_1+2i+2j+k;\gamma+2i+j+k;y\right) \tag{4.14}$$

$$H_C\left(\alpha,\beta_1,\beta_2;\gamma\,;x,y,y\right)=$$

$$=\sum_{i,j,k,l=0}^{\infty}\frac{(-1)^{i+j}\,(\alpha)_{i+2j+k+l}\,(\beta_1)_{2i+j+k+l}\,(\beta_2)_{i+j+k+l}\,(\gamma)_{2i+2j}}{(\gamma+i+j-1)_{i+j}\,(\gamma)_{2i+2j+k+l}\,(\gamma)_{2i+2j+k+l}\,i!j!k!l!}x^{i+j+k+l}y^{i+j+k+l}\cdot$$

$$\cdot F\left(\alpha+i+2j+k+l,\beta_1+2i+j+k+l;\gamma+2i+2j+k+l\,;x\right)$$

$$F\left(\beta_2+i+j+k+l;\alpha+\beta_1+3i+3j+2k+l;\gamma+2i+2j+k+l\,;y\right) \tag{4.15}$$

## 5. Proofs of the received decompositions.

**Case 1.** Decomposition (4.1) can be proved with the help of integral representation, i.e.:

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=\frac{\Gamma\left(\gamma_1\right)\Gamma\left(\gamma_2\right)}{\Gamma\left(\beta_1\right)\Gamma\left(\beta_2\right)\Gamma\left(\gamma_1-\beta_1\right)\Gamma\left(\gamma_2-\beta_2\right)}\cdot$$

$$\cdot\int\limits_0^1\int\limits_0^1\xi^{\beta_1-1}\eta^{\beta_2-1}\left(1-\xi\right)^{\gamma_1-\beta_1-1}\left(1-\eta\right)^{\gamma_2-\beta_2-1}\left(1-y\eta\right)^{\alpha-\beta_1}\left[\left(1-y\eta\right)\left(1-z\eta\right)-x\xi\right]^{-\alpha}d\xi\,d\eta, \tag{5.1}$$

$$\mathrm{Re}\beta_1>0,\ \mathrm{Re}\beta_2>0,\ \mathrm{Re}\left(\gamma_1-\beta_1\right)>0,\ \mathrm{Re}\left(\gamma_2-\beta_2\right)>0.$$

We shall prove decomposition (4.1). For this purpose, we shall take advantage of integral representation (5.1). Taking into account formula

$$\left[\left(1-y\eta\right)\left(1-z\eta\right)-x\xi\right]^{-\alpha}=\left[\left(1-x\xi\right)\left(1-y\eta\right)\left(1-z\eta\right)\right]^{-\alpha}\sum_{i,j=0}^{\infty}\frac{(\alpha)_{i+j}}{i!j!}\sigma_1^i\sigma_2^j, \tag{5.2}$$

where

$$\sigma_1=\frac{xz\xi\eta}{\left(1-x\xi\right)\left(1-y\eta\right)\left(1-z\eta\right)},\quad \sigma_2=\frac{xy\xi\eta}{\left(1-x\xi\right)\left(1-y\eta\right)}. \tag{5.3}$$

Substituting equality (4.2) into integral representation (4.1), we find

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=\sum_{i,j=0}^{\infty}\frac{(\alpha)_{i+j}}{i!j!}x^{i+j}y^jz^i$$

$$\cdot\frac{\Gamma\left(\gamma_1\right)}{\Gamma\left(\beta_1\right)\Gamma\left(\gamma_1-\beta_1\right)}\int\limits_0^1\xi^{\beta_1+i+j-1}\left(1-\xi\right)^{\gamma_1-\beta_1-1}\left(1-x\xi\right)^{-\alpha-i-j}\,d\xi$$

$$\cdot\frac{\Gamma\left(\gamma_2\right)}{\Gamma\left(\beta_2\right)\Gamma\left(\gamma_2-\beta_2\right)}\int\limits_0^1\eta^{\beta_2+i+j-1}\left(1-\eta\right)^{\gamma_2-\beta_2-1}\left(1-y\eta\right)^{-\beta_1-i-j}\left(1-z\eta\right)^{-\alpha-i}d\eta. \tag{5.4}$$

By virtue of integral representations

$$\int\limits_0^1\xi^{b-1}\left(1-\xi\right)^{c-b-1}\left(1-x\xi\right)^{-a}\,d\xi\ =\frac{\Gamma\left(b\right)\Gamma\left(c-b\right)}{\Gamma\left(c\right)}F\left(a,b;c;x\right),$$

$$\mathrm{Re}\,b>0,\ \mathrm{Re}\ (c-b)>0,$$

$$\int\limits_0^1\eta^{a-1}\left(1-\eta\right)^{c-a-1}\left(1-y\eta\right)^{-b_1}\left(1-z\eta\right)^{-b_2}d\eta=\frac{\Gamma\left(a\right)\Gamma\left(c-a\right)}{\Gamma\left(c\right)}F_1\left(a,b_1,b_2;c;y,z\right),$$

$$\mathrm{Re}\,a>0,\ \mathrm{Re}\ (c-a)>0,$$

from (4.4) follows decomposition (3.1).

**Case 2.** The obtained operator identities (2.1) - (2.15) consist of the operator or superposition of operators (1.1) - (1.12). We shall show, as it is possible to apply superpositions of operators for Hypergeometric function. For instance, we consider decomposition (3.7). It's easy to see, that equality takes place

$$\nabla_{xz}(\alpha)\nabla_{xy}(\beta_1)\nabla_{yz}(\beta_2) =$$

$$= \frac{1}{(\beta_2)_n(\beta_2)_p(\alpha)_p}\sum_{i,j,k=0}^{\infty}\frac{(\beta_2)_{n+j}(\beta_2)_{p+i}(\alpha)_{p+i}(-\delta_1)_{i+j}(-\delta_2)_{i+k}(-\delta_3)_{j+k}}{(\alpha)_{i+j}(\beta_1)_i(\beta_2)_{i+j+k}\, i!j!k!}, \tag{5.5}$$

Taking into account the identities (4.5), from parity (2.7), we have

$$H_A(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z) = \sum_{i,j,k=0}^{\infty}\frac{(\beta_2)_j(\beta_2)_i(\alpha)_i(-\delta_1)_{i+j}(-\delta_2)_{i+k}(-\delta_3)_{j+k}}{(\alpha)_{i+j}(\beta_1)_i(\beta_2)_{i+j+k}\, i!j!k!}\cdot$$

$$\cdot F(\alpha,\beta_1;\gamma_1;x)\,F_3(\beta_1,\beta_2+i,\beta_2+j,\alpha+i;\gamma_2;y,z). \tag{5.6}$$

By virtue of the formula [16, p. 93]

$$(\delta+a)(\delta+a+1)\cdots(\delta+a+r-1)f(\xi) = \xi^{1-a}\frac{d^r}{d\xi^{\,r}}[\xi^{a+r-1}f(\xi)], \tag{5.7}$$

where $f(x)$ - analytic function, we find that

$$(-\delta)_r f(\xi) = (-1)^r\xi^r\frac{d^r}{d\xi^{\,r}}f(\xi).$$

We have

$$(-\delta_1)_{i+j}F(\alpha,\beta_1;\gamma_1;x) = (-1)^{i+j}\frac{(\alpha)_{i+j}(\beta_1)_{i+j}}{(\gamma_1)_{i+j}}x^{i+j}F(\alpha+i+j,\beta_1+i+j;\gamma_1+i+j;x), \tag{5.8}$$

$$(-\delta_2)_{i+k}(-\delta_3)_{j+k}F_3(\beta_1,\beta_2+i,\beta_2+j,\alpha+i;\gamma_2;y,z) =$$

$$= (-1)^{i+j}y^{i+k}z^{j+k}\frac{(\alpha)_{i+j+k}(\beta_1)_{i+k}(\beta_2)^2_{i+j+k}}{(\alpha)_i(\beta_1)_j(\beta_2)_i(\gamma_2)_{i+j+2k}}\cdot$$

$$\cdot F_3(\beta_1+i+k,\beta_2+i+j+k,\beta_2+i+j+k,\alpha+i+j+k;\gamma_2+i+j+2k;y,z), \tag{5.9}$$

Substituting identities (4.8) - (4.9) into equality (4.6), we obtain the decomposition (3.7).

**Case 3. The integral representation exists for Hypergeometric function $H_B(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z)$.**

$$H_B(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z) = \frac{\Gamma(\gamma_1)\Gamma(\gamma_2)\Gamma(\gamma_3)}{\Gamma(\beta_1)\Gamma(\beta_2)\Gamma(\alpha)\Gamma(\gamma_1-\beta_1)\Gamma(\gamma_2-\beta_2)\Gamma(\gamma_3-\alpha)}\cdot$$

$$\cdot\int_0^1\int_0^1\int_0^1 \xi^{\beta_1-1}\eta^{\beta_2-1}\zeta^{\alpha-1}(1-\eta)^{\gamma_2-\beta_2-1}(1-x\xi)^{1+\beta_2-\gamma_3}(1-y\eta)^{2+\alpha-\gamma_1-\gamma_3}\cdot$$

$$(1-x\xi-z\zeta)^{1+\beta_1-\beta_2-\gamma_1}\cdot\left[(1-\zeta)(1-x\xi)(1-y\eta)+xy\xi\eta\zeta\right]^{\gamma_3-\alpha-1}\cdot$$

$$\cdot\left[(1-\xi)(1-y\eta)(1-x\xi-z\zeta)+yz\xi\eta\zeta\right]^{\gamma_1-\beta_1-1}d\xi d\eta d\zeta \tag{5.10}$$

$\mathrm{Re}\alpha>0, \mathrm{Re}\beta_1>0,\ \mathrm{Re}\beta_2>0,\ \mathrm{Re}\left(\gamma_1-\beta_1\right)>0,\ \mathrm{Re}\left(\gamma_2-\beta_2\right),\ \mathrm{Re}\left(\gamma_3-\alpha\right)>0.$

We shall prove the decomposition (3.11).

$$\left[(1-\zeta)(1-x\xi)(1-y\eta)+xy\xi\eta\zeta\right]^{\gamma_3-\alpha-1}=(1-x\xi)^{\gamma_3-\alpha-1}(1-y\eta)^{\gamma_3-\alpha-1}(1-\zeta)^{\gamma_3-\alpha-1}\cdot$$

$$\cdot\sum_{i=0}^{\infty}\frac{(1+\alpha-\gamma_3)_i}{i!}\left(-\frac{xy\xi\eta\zeta}{(1-x\xi)(1-y\eta)(1-\zeta)}\right)^i, \tag{5.11}$$

$$\left[(1-\xi)(1-y\eta)(1-x\xi-z\zeta)+yz\xi\eta\zeta\right]^{\gamma_1-\beta_1-1}=(1-\xi)^{\gamma_1-\beta_1-1}(1-y\eta)^{\gamma_1-\beta_1-1}\cdot$$

$$\cdot(1-x\xi-z\zeta)^{\gamma_1-\beta_1-1}\sum_{j=0}^{\infty}\frac{(1+\beta_1-\gamma_1)}{j!}\left(-\frac{yz\xi\eta\zeta}{(1-\xi)(1-y\eta)(1-x\xi-z\zeta)}\right)^j. \tag{5.12}$$

Substituting (4.11)-(4.12) in integral representation (4.10), we define

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)=\frac{\Gamma(\gamma_1)\Gamma(\gamma_2)\Gamma(\gamma_3)}{\Gamma(\beta_1)\Gamma(\beta_2)\Gamma(\alpha)\Gamma(\gamma_1-\beta_1)\Gamma(\gamma_2-\beta_2)\Gamma(\gamma_3-\alpha)}\cdot$$

$$\cdot\sum_{i,j=0}^{\infty}\frac{(-1)^{i+j}(1+\alpha-\gamma_3)_i(1+\beta_1-\gamma_1)_j}{i!j!}x^iy^{i+j}z^j\cdot$$

$$\cdot\int_0^1\int_0^1\int_0^1 \xi^{\beta_1-1+i+j}\eta^{\beta_2-1+i+j}\zeta^{\alpha-1+i+j}(1-\xi)^{\gamma_1-\beta_1-1-j}(1-\eta)^{\gamma_2-\beta_2-1}(1-\zeta)^{\gamma_3-\alpha-1-i}\cdot$$

$$\cdot(1-x\xi)^{\beta_2-\alpha-i}(1-y\eta)^{-\beta_1-i-j}(1-x\xi-z\zeta)^{-\beta_2-j}d\xi d\eta d\zeta. \tag{5.13}$$

By virtue of parity

$$(1-x\xi-z\zeta)^{-\beta_2-j}=(1-x\xi)^{-\beta_2-j}(1-z\zeta)^{-\beta_2-j}\sum_{k=0}^{\infty}\frac{(\beta_2+j)_k}{k!}\left(\frac{xz\xi\zeta}{(1-x\xi)(1-z\zeta)}\right)^k, \tag{5.14}$$

we have

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)$$

$$=\sum_{i,j,k=0}^{\infty}\frac{(-1)^{i+j}(1+\alpha-\gamma_3)_i(1+\beta_1-\gamma_1)_j(\beta_2+j)_k}{i!j!k!}x^{i+k}y^{i+j}z^{j+k}$$

$$\cdot\frac{\Gamma(\gamma_1)}{\Gamma(\beta_1)\Gamma(\gamma_1-\beta_1)}\int_0^1 \xi^{\beta_1-1+i+j+k}(1-\xi)^{\gamma_1-\beta_1-1-j}(1-x\xi)^{-\alpha-i-j-k}d\xi$$

$$\cdot\frac{\Gamma(\gamma_2)}{\Gamma(\beta_2)\Gamma(\gamma_2-\beta_2)}\int\limits_0^1 \eta^{\beta_2-1+i+j}(1-\eta)^{\gamma_2-\beta_2-1}(1-y\eta)^{-\beta_1-i-j}\,d\eta$$

$$\cdot\frac{\Gamma(\gamma_3)}{\Gamma(\alpha)\Gamma(\gamma_3-\alpha)}\int\limits_0^1 \zeta^{\alpha-1+i+j+k}(1-\zeta)^{\gamma_3-\alpha-1-i}(1-z\zeta)^{-\beta_2-j-k}\,d\zeta. \tag{5.15}$$

and from identity (5.15) we define the decomposition (3.11).

**Case 4.** Decomposition (3.8) can be proved by means of equality

$$\nabla_{xz}(\alpha)\nabla_{xy}(\beta_1)\nabla_{yz}(\beta_2)\nabla_{yz}(\gamma)=\frac{1}{(\alpha)_m(\alpha)_p(\beta_1)_m(\beta_2)_n(\beta_2)_p}$$

$$\cdot\sum_{i,j,k,l=0}^{\infty}\frac{(\alpha)_{j+k}(\alpha)_k(\beta_1)_{i+j}(\beta_2)_i^2(\alpha+j+k)_m(\alpha+k)_p(\beta_1+i+j)_m(\beta_2+i)_n(\beta_2+i)_p}{(\alpha)_{i+j+k+l}(\beta_1)_{i+j+k}(\beta_2)_{2i+j}(\gamma)_i}$$

$$\cdot\frac{(-\delta_1)_{k+l}(-\delta_2)_{i+j+k}(-\delta_3)_{i+j+l}}{i!j!k!l!} \tag{5.16}$$

Taking into account the identities (4.16), from parity (2.8), we have

$$H_B(\alpha,\beta_1,\beta_2;\gamma_1,\gamma,\gamma;x,y,z)=\sum_{i,j,k,l=0}^{\infty}\frac{(\alpha)_{j+k}(\alpha)_k(\beta_1)_{i+j}(\beta_2)_i^2}{(\alpha)_{i+j+k+l}(\beta_1)_{i+j+k}(\beta_2)_{2i+j}(\gamma)_i\,i!j!k!l!}\cdot$$

$$\cdot(-\delta_1)_{k+l}F(\alpha+j+k,\beta_1+i+j;\gamma_1;x)\cdot$$

$$\cdot(-\delta_2)_{i+j+k}(-\delta_3)_{i+j+l}F_3(\beta_2+i,\alpha+k,\beta_1,\beta_2+i;\gamma;y,z) \tag{5.17}$$

We have

$$(-\delta_1)_{k+l}F(\alpha+j+k,\beta_1+i+j;\gamma_1;x)=$$

$$=(-1)^{k+l}x^{k+l}\frac{(\alpha)_{j+2k+l}(\beta_1)_{i+j+k+l}}{(\alpha)_{j+k}(\beta_1)_{i+j}(\gamma_1)_{k+l}}F(\alpha+j+2k+l,\beta_1+i+j+k+l;\gamma_1+k+l;x), \tag{5.18}$$

$$(-\delta_2)_{i+j+k}(-\delta_3)_{i+j+l}F_3(\beta_2+i,\alpha+k,\beta_1,\beta_2+i;\gamma;y,z)=$$

$$=(-1)^{k+l}y^{i+j+k}z^{i+j+l}\frac{(\alpha)_{i+j+k+l}(\beta_1)_{i+j+k}(\beta_2)_{2i+j+k}(\beta_2)_{2i+j+l}}{(\alpha)_k(\beta_2)_i^2(\gamma)_{2i+2j+k+l}}$$

$$F_3(\beta_2+2i+j+k,\alpha+i+j+k+l,\beta_1+i+j+k,\beta_2+2i+j+l;$$

$$\gamma+2i+2j+k+l;y,z) \tag{5.19}$$

Substituting identities (4.18) - (4.19) into equality (4.17), we obtain the decomposition (3.8).

## 6. The Integrals.

With the help of the obtained decompositions can be easily found the following integrals connecting with Hypergeometric $H_A,\ H_B$ functions

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=\frac{\Gamma\left(\gamma_1\right)\Gamma\left(\gamma_2\right)}{\Gamma\left(\beta_1\right)\Gamma\left(\beta_2\right)\Gamma\left(\gamma_1-\beta_1\right)\Gamma\left(\gamma_2-\beta_2\right)}\cdot$$

$$\cdot\int\limits_0^1\int\limits_0^1\xi^{\beta_1-1}\eta^{\beta_2-1}\left(1-\xi\right)^{\gamma_1-\beta_1-1}\left(1-\eta\right)^{\gamma_2-\beta_2-1}\left(1-y\eta\right)^{-\beta_2}\left(1-x\xi-z\eta\right)^{-\alpha}\cdot$$

$$\cdot F\left(\alpha,\beta_2;\beta_1;\frac{xy\xi\eta}{\left(1-y\eta\right)\left(1-x\xi-z\eta\right)}\right)d\xi\,d\eta,\tag{6.1}$$

$$\mathrm{Re}\alpha_1>0,\quad\mathrm{Re}\left(\gamma_1-\alpha_1\right)>0,\quad\ \mathrm{Re}\alpha_2>0,\quad\mathrm{Re}\left(\gamma_2-\alpha_2\right)>0,$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2,\gamma_3;x,y,z\right)=\frac{\Gamma\left(\gamma_1\right)\Gamma\left(\gamma_2\right)\Gamma\left(\gamma_3\right)}{\Gamma^2\left(\alpha\right)\Gamma\left(\beta_1\right)\Gamma\left(\gamma_1-\alpha\right)\Gamma\left(\gamma_2-\beta_1\right)\Gamma\left(\gamma_3-\alpha\right)}\cdot$$

$$\cdot\int\limits_0^1\int\limits_0^1\int\limits_0^1\xi^{\alpha-1}\eta^{\beta_1-1}\zeta^{\alpha-1}\left(1-\xi\right)^{\gamma_1-\alpha-1}\left(1-\eta\right)^{\gamma_2-\beta_1-1}\left(1-\zeta\right)^{\gamma_3-\alpha-1}\left(1-x\xi\right)^{\beta_2-\beta_1}\cdot$$

$$\cdot\left[\left(1-x\xi\right)\left(1-y\eta-z\zeta\right)-xy\xi\eta\right]^{-\beta_2}\cdot$$

$$\cdot F\left(\beta_2,1+\beta_1-\gamma_2;\alpha;-\frac{xz\xi\eta\zeta}{\left(1-\eta\right)\left[\left(1-x\xi\right)\left(1-y\eta-z\zeta\right)-xy\xi\eta\right]}\right)d\xi d\eta d\zeta,\tag{6.2}$$

$$\mathrm{Re}\alpha>0,\quad\mathrm{Re}\left(\gamma_1-\alpha\right)>0,\ \mathrm{Re}\beta_1>0,\quad\mathrm{Re}\left(\gamma_2-\beta_1\right)>0,\quad\mathrm{Re}\left(\gamma_3-\alpha\right)>0.$$

These identities can be proved by using the expansions of under integral functions.

If we use the definition for Euler function $\Gamma\left(z\right)$ ([13], Ch. 1, Sec. 1.2, (1))

$$\Gamma\left(z\right)=\left\{\begin{array}{c}\int\limits_0^\infty e^{-t}t^{z-1}dt,\ \mathrm{Re}\ (z)>0,\\ \frac{\Gamma(1+z)}{z},\mathrm{Re}\ (z)<0;\ z\neq-1,-2,....\end{array}\right.\tag{6.3}$$

Then, for Hypergeometric $H_A$, $H_B$, $H_C$ functions we find the following integral representations

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)=\frac{1}{\Gamma\left(\alpha\right)\Gamma\left(\beta_1\right)\Gamma\left(\beta_2\right)}$$

$$\cdot\int\limits_0^\infty\int\limits_0^\infty\int\limits_0^\infty e^{-u_1-u_2-u_3}u_1^{\alpha-1}u_2^{\beta_1-1}u_3^{\beta_2-1}{}_0F_1\left(\gamma_1;xu_1u_2\right){}_0F_1\left(\gamma_2;yu_2u_3+zu_1u_3\right)du_1du_2du_3,\tag{6.4}$$

$$\mathrm{Re}\,\alpha>0,\ \mathrm{Re}\,\beta_1>0,\ \mathrm{Re}\,\beta_2>0,$$

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\gamma_2;x,y,z\right)$$

$$= \frac{1}{\Gamma(\alpha)\Gamma(\beta_1)} \int_0^\infty \int_0^\infty e^{-u_1-u_2} u_1^{\alpha-1} u_2^{\beta_1-1} {}_0F_1(\gamma_1; xu_1u_2)\ {}_1F_1(\beta_2; \gamma_2; yu_2 + zu_1)\, du_1 du_2, \quad (6.5)$$

$$\operatorname{Re}\alpha > 0,\ \operatorname{Re}\beta_1 > 0,$$

$$H_B(\alpha, \beta_1, \beta_2; \gamma_1, \gamma_2, \gamma_3; x, y, z) = \frac{1}{\Gamma(\alpha)\Gamma(\beta_1)\Gamma(\beta_2)}$$

$$\cdot \int_0^\infty \int_0^\infty \int_0^\infty e^{-\xi-\eta-\zeta} \xi^{\alpha-1} \eta^{\beta_1-1} \zeta^{\beta_2-1} {}_0F_1(\gamma_1; x\xi\eta)\ {}_0F_1(\gamma_2; y\eta\zeta)\ {}_0F_1(\gamma_3; z\xi\zeta)\, d\xi d\eta d\zeta, \quad (6.6)$$

$$\operatorname{Re}\alpha > 0,\ \operatorname{Re}\beta_1 > 0,\ \operatorname{Re}\beta_2 > 0,$$

$$H_B(\alpha, \beta_1, \beta_2; \gamma_1, \gamma_2, \gamma_3; x, y, z)$$

$$= \frac{1}{\Gamma(\alpha)\Gamma(\beta_1)} \int_0^\infty \int_0^\infty e^{-\xi-\eta} \xi^{\alpha-1} \eta^{\beta_1-1} {}_0F_1(\gamma_1; x\xi\eta)\, \Psi_2(\beta_2; \gamma_2, \gamma_3; y\eta, z\xi)\, d\xi d\eta, \quad (6.7)$$

$$\operatorname{Re}\alpha > 0,\ \operatorname{Re}\beta_1 > 0,$$

$$H_C(\alpha, \beta_1, \beta_2; \gamma; x, y, z)$$

$$= \frac{1}{\Gamma(\alpha)\Gamma(\beta_1)\Gamma(\beta_2)} \int_0^\infty \int_0^\infty \int_0^\infty e^{-\xi-\eta-\zeta} \xi^{\alpha-1} \eta^{\beta_1-1} \zeta^{\beta_2-1} {}_0F_1(\gamma; x\xi\eta + y\eta\zeta + z\xi\zeta)\, d\xi d\eta d\zeta, \quad (6.8)$$

$$\operatorname{Re}\alpha > 0,\ \operatorname{Re}\beta_1 > 0,\ \operatorname{Re}\beta_2 > 0,$$

where [15]

$$\Psi_2(\alpha; \gamma_1, \gamma_2; x, y) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_{m+n}}{(\gamma_1)_m (\gamma_2)_n m! n!} x^m y^n, \quad (6.9)$$

and, as usual ${}_pF_q$ denotes the generalized Hypergeometric function with $p$ numerator and $q$ denominator parameters. For multiple hypergeometric of functions $H_B^{(n)}$, $H_C^{(n)}$ the integrated representations are determined in the monograph ([13], Ch. 9, Sec. 9.4, (198)-(200)).

## 7. Conclusion.

From decomposions (4.1)-(4.15) as a private value can be found the known decompositions that have been found in [5], i.e. decompositions (4.1)-(4.15) generalize the results of [5]. Further, the following decompositions for Hypergeometric Appell's functions

$$F_1(\alpha; \beta, \beta; \gamma; x, y) = \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j} (\alpha)_{2i+2j} (\beta)_i (\beta)_{i+j} (\gamma)_{2i}}{(\gamma + i - 1)_i (\gamma)_{2i+j}^2 i! j!} x^{i+j} y^{i+j}$$

$$F_4(\alpha + 2i + 2j, \beta + i + j; \gamma + 2i + j, \gamma + 2i + j; x, y), \quad (7.1)$$

$$F_2\left(\alpha;\beta_1,\beta_2;\gamma,\gamma;x,y\right)=\sum_{i,j=0}^{\infty}\frac{(\alpha)_{2i+j}\,(\beta_1)_{i+j}\,(\beta_2)_{i+j}}{(\gamma)_i\,(\gamma)_{2i+2j}\,i!j!}x^{i+j}y^{i+j}$$
$$F_3\left(\alpha+2i+j,\alpha+2i+j;\beta_1+i+j,\beta_2+i+j;\gamma+2i+2j;x,y\right), \tag{7.2}$$

have not been found in work [5]. In particular, from decompositions (4.1), (4.2) we can find the following identities

$$H_A\left(\alpha,\beta_1,\beta_2;\gamma_1,\beta_2;x,y,z\right)=(1-y)^{-\beta_1}(1-z)^{-\alpha}F\left(\alpha,\beta_1;\gamma_1;\frac{x}{(1-y)(1-z)}\right), \tag{7.3}$$

$$H_B\left(\alpha,\beta_1,\beta_2;\gamma_1,\beta_2,\beta_2;x,y,z\right)$$
$$=(1-y)^{-\beta_1}(1-z)^{-\alpha}F_4\left(\alpha,\beta_1;\beta_2,\gamma_1;\frac{yz}{(1-y)(1-z)},\frac{x}{(1-y)(1-z)}\right). \tag{7.4}$$

It should be noted that the following superpositions of the operators

$$\nabla_{xy}(\alpha)\nabla_{xz}(\alpha)\Delta_{yz}(\alpha)\Delta_{yz}(\gamma_2), \quad \nabla_{xz}(\alpha)\nabla_{xy}(\alpha)\Delta_{yz}(\alpha),$$

$$\nabla_{xz}(\alpha)\nabla_{xy}(\alpha)\Delta_{yz}(\alpha)\Delta_{yz}(\gamma)\tilde{\Delta}_x(\gamma)$$

have not been applied for Hypergeometric functions.

## REFERENCES

[1] G.Lohofer. Theory of an electromagnetically levitated metal spheres. Absorbed Power. SIAM J. Appl. Math., 1989, 49, N2, 567-581.

[2] A.W.Ninkkanen. Generalized hypergeometric series ${}^{N}F\left(x_1,x_2,...,x_N\right)$ , arising in physical and quantum chemical applications. J. Phys. A: Math. and Gen., 1983, 16, N9, 1813-1825.

[3] F. I. Frankl. The elected works on gas dynamics. Moscow, 1973.

[4] A. Hasanov. About one mixed task for the equation $signy\,|y|^m\,u_{xx}+x^n u_{yy}=0$. New. Academ. Scien. Uz SSR, N 2, 1982, 28-32.

[5] J. L. Burchnall and T.W. Chaundy. Expansions of Appell's double hypergeometric functions. Quart. J. Math., Oxford, 1940, Ser. 11, 249-270.

[6] J. L. Burchnall and T.W. Chaundy. Expansions of Appell's double hypergeometric functions (II). Quart. J. Math., Oxford, 1941, Ser. 12, 112-128.

[7] T.W. Chaundy. Expansions of hypergeometric functions. Quart. J. Math., Oxford, 1942, Ser. 13, 159-171.

[8] Srivastava H.M., Hypergeometric functions of three variables. Ganita, 1964, 15, N2, 97-108.

[9] J. P. Singhal and S. S. Bhati, Certain expansions associated with hypergeometric functions of $n$ variables, Glasnik Mat. Ser. 3, 11 (31), 1976, 239-245.

[10] H. M. Srivastava, Some integrals representing triple hypergeometric functions, Rend. Circ. Mat. Palermo, (ser. 2) 16, 1967, 99-115.

[11] G. Lauricella, Sulle funzioni ipergeometriche a piu variabili, Rend. Circ. Mat. Palermo 7, (1893), 111-158.

[12] P. Appell and J. Kampe de Feriet, Fonctions hypergeometriques et hyperspheriques. Polynomes d'Hermite, Gauthier - Villars. Paris, 1926.

[13] H. M. Srivastava and P. W. Karlsson, Multiple Gaussian Hypergeometric Series. Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.

[14] O. I. Marichev. Handbook of integral transforms of higher transcendental functions, theory and algorithmic tables. Chichester, Ellis Horwood, 1982.

[15] A. Erdelyi (Ed): Higher transcendental functions, vol. 1. New York: McGraw Hill Book Company, Inc. 1953.

[16] E. G. C. Poole. Introduction to the theory of linear differential equation. Oxford, 1936.